SURGICAL GROUP (941) 896-4788	Patient Info: Post-Op Instructions for Gallbladder Surgery by Dr. Jose Erbella
Procedure	Laparoscopic Cholecystectomy
Incision Care	Remove any dressing(s) in 24hrs. The incision(s) may be sealed with staples , a skin adhesive (Dermabond) or covered with white adhesive tapes known as steri-strips . Either way, there is no need to cover up the incisions again with gauze.
Shower	You may shower 24 hours after surgery & get the incision(s) or steri-strips wet with soapy water and gently pad them dry. If the steri-strips come off in the shower, do not become concerned. If they are still in place 10 days after surgery, you may remove them.
Pain Medication	You will be given a prescription for a narcotic pain medication (usually Lortab or Percocet). Options for non-constipating pain medications include products that include Ibuprofen or Tylenol. To combat constipation, you may take a stool softener (Colace) twice each day.
Common Complaints	Shoulder pain from the carbon dioxide used during the procedure is common. Your body absorbs this gas naturally over a 48-72 hour period. Tylenol (Acetaminophen) or Ibuprofen (Motrin) is usually sufficient to relieve this discomfort. Your throat may be sore from the breathing tube. This too resolves in a day or two. Cough drops and throat lozenges are soothing.
Activity	Do not do any heavy lifting (more than 10 pounds) or vigorous exercise for 4-6 weeks .
Diet	Stick to a low-fat diet for at least the first few weeks. Fatty foods such as fried chicken or a greasy cheeseburger may lead to diarrhea. If your throat is sore, try eating soft foods.
Return to Work	You may return to work in 1-2 weeks .
Physician Appointment	Call (941) 896-4788 to make your post-operative appointment to see Dr. Erbella in 1-2 weeks or if you have any questions.
When to call your Surgeon	Persistent fever over 101.5 degrees F (39 C). Pain that is not relieved by your medications. Persistent nausea or vomiting. Persistent cough or shortness of breath. Purulent drainage (pus) or bleeding from any incision. Redness surrounding any of your incisions that is worsening or getting bigger. If you are unable to eat or drink liquids.
In Case of Emergency	Dr. Erbella is available 24hrs a day. You may call either our office (941) 896-4788, your nearest emergency room at Blake, Lakewood Ranch or Manatee Hospitals or Dial 911.